

1-phenyl-2-propanone

AI generated article from Bing

$$\square\square - \square\square$$
[illegible]

10 - 10

□□□□□□□□ 1+1=2□□□□□□ 1 □□□□□□ 0 □□□□□□□□ 0 □

000 **1** 0000000000 - 00

[illegible]

1-1+1-1+1-1+1... □□□□□□□□□□□□□□□□ - □□

2011 1

Formal proof for $(-1) \times (-1) = 1$ - Mathematics Stack Exchange

Is there a formal proof for $(-1) \times (-1) = 1$? It's a fundamental formula not only in arithmetic but also in the whole of math. Is there a proof for it or is it just assumed?

$$1/1+1/2+1/3+1/4+.....+1/n=\square\square\square\square\square\square - \square\square$$
$$\frac{1}{n} \left(\ln n - \ln \left(\frac{n-1}{n} \right) \right) = \frac{1}{n} \left(\ln n - \ln (n-1) + \ln n \right) = \frac{1}{n} \left(2 \ln n - \ln (n-1) \right)$$

1/8, 1/4, 1/2, 3/4, 7/8 □□□□□□□□? - □□

1/8 1/4 3/8 1/2 5/8 3/4 7/8 This is an arithmetic sequence since there is a common difference between each term. In this case, adding 1/8 to the previous term in the sequence gives the next term. In other words, $a_n = a_1 + d(n-1)$. Arithmetic Sequence: $d=1/8$

□ □ - □ □ □ □ □ □ □

2011 1

Word 1.1 2.1 1.1 ? -

1 2 3 —

What is the value of 1^i ? - Mathematics Stack Exchange

The formal moral of that example is that the value of 1^i depends on the branch of the complex logarithm that you use to compute the power. You may already know that $1 = e^{0+2k\pi i} = e^{0 + 2k\pi i}$ for every integer k , so there are many possible choices for $\log(1)$.