

dod directive 5240.06

AI generated article from Bing

Priapism - Diagnosis & treatment - Mayo Clinic

Diagnostic tests Diagnostic tests might be needed to determine what type of priapism you have. Additional tests might identify the cause of priapism. In an emergency room setting, your treatment will likely begin before all test results are received. Diagnostic tests can include: Penile blood gas measurement. In this test, a tiny needle is inserted into your penis to remove a sample of blood ...

Victoria E. Clark, M.D., Ph.D. - Mayo Clinic

Victoria E. Clark, M.D., Ph.D., is a Senior Associate Consultant in the Department of Neurologic Surgery and the State of Florida Endowed Chair of Cancer Research at the Mayo Clinic's Florida campus. Dr. Clark graduated with a BA with high honors in Biochemical Sciences from Harvard. She earned her MD and PhD degrees from the Yale University School of Medicine, where she was awarded the degree ...

Animal bites: First aid - Mayo Clinic

Animal bites might be caused by pets — such as cats, dogs, hamsters and turtles. Or bites may be from farm animals or wild animals. The type of care needed depends on how deep the bite is and the type of animal that caused it. You may need medicine to fight infection. Or you may need a tetanus ...

Pemphigus - Symptoms and causes - Mayo Clinic

Learn about symptoms, causes, treatment and self-care tips for this group of rare conditions that causes blisters and sores on the skin or mucous membranes.

Stye (sty) - Diagnosis & treatment - Mayo Clinic

In most cases, a stye doesn't require specific treatment, but using warm compresses can hasten the healing. A stye typically goes away on its own. Recurrences are common. For a stye that won't go away, additional treatment may be recommended, such as: Antibiotics. A healthcare professional may prescribe antibiotic eye drops or a topical antibiotic cream to apply to the eyelid. If the eyelid ...

Optimizing care for adults with traumatic brain injury

Classifying TBI severity Many medical centers use the VA/DoD Clinical Practice Guideline for the Management and Rehabilitation of Post-Acute Mild Traumatic Brain Injury (mTBI), based on the Glasgow Coma Scale (GCS), loss of consciousness duration and post-traumatic amnesia.

Black eye: First aid - Mayo Clinic

A black eye is bruising caused by bleeding in the tiny blood vessels in the skin surrounding the eye. Most injuries that cause a black eye aren't serious. But a black eye could be a symptom of a more

serious injury, such as an internal injury to the eye or a fracture of the thin bones around the eye ...

Oppositional defiant disorder (ODD) - Symptoms and causes

Overview Even the best-behaved children can be difficult and challenging at times. But oppositional defiant disorder (ODD) includes a frequent and ongoing pattern of anger, irritability, arguing and defiance toward parents and other authority figures. ODD also includes being spiteful and seeking revenge, a behavior called vindictiveness.

Blood in urine (hematuria) - Symptoms and causes - Mayo Clinic

Symptoms Blood in the urine can look pink, red or cola-colored. Red blood cells cause the urine to change color. It takes only a small amount of blood to turn urine red. The bleeding often isn't painful. But if blood clots get passed in the urine, that can hurt. See a health care provider whenever urine looks like it might have blood in it. Red urine isn't always caused by red blood cells ...

Anxiety disorders - Diagnosis and treatment - Mayo Clinic

Diagnosis You may start by seeing your primary care provider to find out if your anxiety could be related to your physical health. He or she can check for signs of an underlying medical condition that may need treatment. However, you may need to see a mental health specialist if you have severe anxiety. A psychiatrist is a medical doctor who specializes in diagnosing and treating mental health ...