

doa tahajud

AI generated article from Bing

Tuberculosis Elimination - Mayo Clinic

a person is infected with *Mycobacterium tuberculosis*. Reliable administration and reading of the TST requires standardization of procedures, training, supervision, and practice.

Needle biopsy - Mayo Clinic

A needle biopsy is a procedure to remove some cells or a small piece of tissue from the body using a needle. The sample removed during a needle biopsy goes to a lab for testing. Common needle biopsy procedures include fine-needle aspiration and core needle biopsy. Needle biopsy may be used to take tissue or fluid samples from the lymph nodes, liver, lungs or bones. It also may be used on other

...

Metoprolol (intravenous route) - Side effects & uses - Mayo Clinic

Description Metoprolol injection is used to reduce the risk of death from an acute heart attack. It is given to people who have already had a heart attack. This medicine is a beta-blocker. It works by affecting the response to nerve impulses in certain parts of the body, like the heart. As a result, the heart beats slower and decreases the blood pressure. When the blood pressure is lowered ...

Electrocardiogram (ECG or EKG) - Mayo Clinic

An electrocardiogram (ECG or EKG) is a quick test to check the heartbeat. It records the electrical signals in the heart. Test results can help diagnose heart attacks and irregular heartbeats, called arrhythmias. ECG machines can be found in medical offices, hospitals, operating rooms and ambulances ...

ANA test - Mayo Clinic

An antinuclear antibody (ANA) test is one test used to help determine if you have an autoimmune disease such as lupus.

Ibuprofen (oral route) - Side effects & dosage - Mayo Clinic

Description Ibuprofen is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, and helps to relieve symptoms of arthritis (osteoarthritis, rheumatoid arthritis, or juvenile arthritis), such as inflammation, swelling, stiffness, and joint pain. This medicine does not cure arthritis and will help you only as long as you continue to take it . In addition, ibuprofen ...

Tricyclic antidepressants - Mayo Clinic

Tricyclic antidepressants can have more side effects than other antidepressants. But for some people, they may ease depression when other medicines fail.

Do drug-eluting stents increase heart attack risk? - Mayo Clinic

A stent is a small mesh tube put into an artery to keep it open. A drug-eluting stent is coated with a slow-release medicine to help stop blood clots from forming in a stent. Blood clotting in a stent can cause a future blockage, called restenosis, and may lead to a heart attack. Stents without a ...

HIDA scan - Mayo Clinic

Overview A hepatobiliary iminodiacetic acid (HIDA) scan is an imaging procedure used to diagnose problems of the liver, gallbladder and bile ducts. For a HIDA scan, also known as cholescintigraphy or hepatobiliary scintigraphy, a radioactive tracer is injected into a vein in the arm. The tracer travels through the bloodstream to the liver, where the bile-producing cells take it up. The tracer ...

5K run: 7-week training schedule for beginners - Mayo Clinic

Think about using this seven-week 5K run training schedule as your guide. It's made for beginners or anyone who wants to do a 5K race. You also can adapt it for a 5K walk. This 5K training schedule includes a mix of running, walking and resting. This mix helps lower the risk of injury, stress and fatigue. It also can help you enjoy physical activity. Keep in mind, you can run or walk slowly to ...